

GATA-dependent recruitment of MEF2 proteins to target promoters

Steves Morin¹, Frédéric Charron^{1,2},
Lynda Robitaille¹ and Mona Nemer^{1,2,3,4}

¹Laboratoire de développement et différenciation cardiaques, Institut de recherches cliniques de Montréal, 110 des Pins Ouest, Montréal, QC, H2W 1R7, ²Department of Medicine, Division of Experimental Medicine, McGill University, Montréal, QC and ³Département de Pharmacologie, Université de Montréal, QC, Canada

⁴Corresponding author at: Institut de recherches cliniques de Montréal, Laboratoire de développement et différenciation cardiaques, 110 des Pins Ouest, Montréal, QC, Canada H2W 1R7
e-mail: nemem@ircm.qc.ca

S.Morin and F.Charron contributed equally to this work

The myocyte enhancer factor-2 (MEF2) proteins are MADS-box transcription factors that are essential for differentiation of all muscle lineages but their mechanisms of action remain largely undefined. In mammals, the earliest site of MEF2 expression is the heart where the MEF2C isoform is detectable as early as embryonic day 7.5. Inactivation of the *MEF2C* gene causes cardiac developmental arrest and severe down-regulation of a number of cardiac markers including atrial natriuretic factor (ANF). However, most of these promoters contain no or low affinity MEF2 binding sites and they are not significantly activated by any MEF2 proteins in heterologous cells suggesting a dependence on a cardiac-enriched cofactor for MEF2 action. We provide evidence that MEF2 proteins are recruited to target promoters by the cell-specific GATA transcription factors, and that MEF2 potentiates the transcriptional activity of this family of tissue-restricted zinc finger proteins. Functional MEF2/GATA-4 synergy involves physical interaction between the MEF2 DNA-binding domain and the carboxy zinc finger of GATA-4 and requires the activation domains of both proteins. However, neither MEF2 binding sites nor MEF2 DNA binding capacity are required for transcriptional synergy. The results unravel a novel pathway for transcriptional regulation by MEF2 and provide a molecular paradigm for elucidating the mechanisms of action of MEF2 in muscle and non-muscle cells.

Keywords: GATA factors/heart development/MEF2 proteins/muscle transcription

Introduction

Members of the myocyte enhancer factor-2 (MEF2) family of MADS (MCM1, Agamous, Deficiens, Serum response factor)-box transcription factors are evolutionarily conserved proteins that are expressed at high levels in all muscle cells. MEF2 proteins are also found in non-muscle cells including brain and lymphoid tissue (reviewed in

Black and Olson, 1998). In mammals, the MEF2 family is composed of four members, MEF2A, MEF2B, MEF2C and MEF2D, which form homo- and heterodimers that bind the consensus DNA sequence (T/C)TA(A/T)₄TA(G/A) present in many muscle and non-muscle promoters. MEF2 proteins contain a conserved N-terminal 56 amino acid MADS domain and an adjacent 29 amino acid MEF2 domain, which together mediate DNA binding and dimerization.

Genetic studies have provided evidence for an essential role of MEF2 proteins in muscle-specific gene expression and differentiation of all three muscle lineages. In *Drosophila*, mutation of the *D-mef2* gene results in embryos lacking differentiated skeletal, cardiac and visceral muscle cells (Bour *et al.*, 1995; Lilly *et al.*, 1995; Ranganayakulu *et al.*, 1995). In mice, inactivation of the *MEF2C* gene, which is the first MEF2 isoform expressed during embryonic development, leads to cardiac morphogenetic defects, vascular abnormalities and lethality by embryonic day 9.5 (Lin *et al.*, 1997, 1998; Bi *et al.*, 1999). The mechanisms by which MEF2 proteins regulate myogenesis of both striated and smooth muscle cells and the identity of their downstream targets in these various tissues are only starting to be elucidated. At present, the mechanisms of action of MEF2 have been analyzed mostly in skeletal muscle where MEF2 appears to act as cofactors for the myogenic basic helix-loop-helix (bHLH) proteins, MyoD, Myf5, myogenin and MRF4 (Kaushal *et al.*, 1994; Molkentin *et al.*, 1995; Black *et al.*, 1998). Thus, MEF2 proteins strongly potentiate the transcriptional activity of the bHLH myogenic factors and cooperate with them for inducing and maintaining the skeletal muscle phenotype. This cooperativity is mediated by direct interaction between the DNA-binding domains of MEF2 and myogenic proteins, and necessitates a DNA-binding site for only one of the two factors. Therefore, in skeletal myocytes, MEF2 may modulate transcription by two distinct pathways: one involving DNA binding to MEF2 sites and another one involving recruitment of MEF2 to E-boxes in target promoters via the myogenic bHLH factors. Whether similar mechanisms underlie the action of MEF2 in cardiac and visceral muscle cells where the MyoD family of transcription factors is not expressed remains unknown.

MEF2-binding sites have been reported in several cardiac promoters and their mutation was shown to decrease promoter activity in cardiomyocytes; they include the MEF2 sites in the ventricular myosin light chain (MLC2V), cardiac troponin T, cardiac troponin I, α -myosin heavy chain (α MHC) and Desmin (Iannello *et al.*, 1991; Zhu *et al.*, 1991; Yu *et al.*, 1992; Molkentin and Markham, 1993; Kuisk *et al.*, 1996; Di Lisi *et al.*, 1998). Analysis of cardiac gene expression in mice with targeted mutation of the *MEF2C* gene confirmed that some of these

genes, like α MHC, required MEF2C for optimum transcription (Lin *et al.*, 1997). In addition to α MHC, two other cardiac-specific genes not previously associated with MEF2 proteins, atrial natriuretic factor (ANF) and α -cardiac actin (α -CA), were completely absent in the hearts of MEF2C-deficient embryos. How MEF2C regulates transcription of these target genes remains unclear; the two α MHC MEF2 sites are low-affinity MEF2-binding sites (Yu *et al.*, 1992; Molkentin and Markham, 1993), and MEF2 proteins are unable to activate α MHC-driven reporters in cotransfection assays although they can potentiate transactivation of the α MHC promoter by the thyroid hormone receptor (Lee *et al.*, 1997). Moreover, ectopic expression of MEF2 proteins in explanted *Xenopus* ectoderm failed to activate endogenous α MHC or α -CA genes (Chambers *et al.*, 1994; Fu and Izumo, 1995). However, forced expression of MEF2 proteins in whole *Xenopus* embryos results in precocious expression of endogenous α MHC and enlarged hearts (Fu and Izumo, 1995). Together, these studies suggest that MEF2 regulates transcription of α MHC and possibly other cardiac genes in conjunction with a cell-specific cofactor present in embryonic mesoderm (or endoderm) but not in ectoderm.

The ANF promoter, a known downstream target for the cardiac-specific transcription factors GATA-4 and Nkx2-5, does not contain MEF2 consensus binding sites (Durocher *et al.*, 1996, 1997; Charron *et al.*, 1999); this suggests an indirect action of MEF2 on ANF transcription possibly through modulation of GATA-4, Nkx2-5 or other ANF regulators. However, neither GATA-4 nor Nkx2-5 levels are altered in MEF2C^{-/-} embryos (Lin *et al.*, 1997).

In this study, we provide evidence that MEF2 proteins are recruited by the cardiac-specific transcription factor GATA-4 to synergistically activate ANF and several other MEF2C target promoters including α MHC and α -CA. The MEF2-GATA-4 synergy is mediated by physical interaction between the respective DNA-binding domains and requires the transactivation domains of both factors. GATA-binding sites are necessary and sufficient for cooperativity with MEF2. Other GATA factors, including GATA-6, which is expressed in cardiac and smooth muscle cells, and GATA-2 and -3, which are present in hemopoietic and neuronal cells, are also able to cooperate with MEF2 proteins. Together, the data suggest that in addition to cooperating with the myogenic bHLH proteins in skeletal muscle differentiation, MEF2 proteins act as cofactors for the tissue-restricted zinc finger GATA proteins in cardiac myogenesis and raise the possibility that GATA factors may be essential components of MEF2 action in several other cell types.

Results

MEF2 proteins activate the ANF promoter via two distinct mechanisms

The absence of ANF transcripts in the heart of mice homozygous for a null mutation of the MEF2C gene (Lin *et al.*, 1997) indicates that ANF is a downstream target for MEF2 proteins. We tested whether the effect of MEF2C was due to direct action on the ANF promoter or via DNA binding or through recruitment by protein-protein interactions. The first 700 bp of the rat ANF promoter are




Fig. 1. The ANF promoter harbors a low-affinity MEF2-binding site. (A) Schematic representation of the ANF promoter. Regulatory elements are boxed and their location relative to the transcription start site is indicated. All these elements are evolutionarily conserved on the ANF promoter from many species. SRE-like is a low-affinity serum response element; the GATA4 and GATA6 are the distal and proximal GATA-binding sites, respectively. The consensus MEF2-binding site is also shown. The A/T-rich mut sequence indicates the mutations introduced to abolish the A/T-rich element. rANF and hANF are the rat and human ANF promoter, respectively. (B) The A/T-rich element is a low-affinity MEF2-binding site. EMSAs were performed on the MEF2 element of the MCK promoter (MEF2-MCK, left panel) or the A/T-rich element of ANF (A/T-rich, right panel) using *in vitro* translated MEF2A. In the left panel, the MEF2A binding was competed with different unlabeled ANF probes described in Materials and methods. Only the A/T-rich element of the ANF promoter was able to compete the MEF2A binding. Similar results were obtained with *in vitro* translated MEF2C and MEF2D.

sufficient to recapitulate cardiac specificity and spatio-temporal regulation of the endogenous gene in cultured cardiomyocytes (Argentin *et al.*, 1994) and in transgenic mice (Durocher *et al.*, 1998).

Sequence analysis of the entire 700 bp rat ANF promoter revealed no consensus MEF2 sites; the closest sequence homologies to MEF2-binding sites mapped to an A/T-rich element sharing similarities with a MEF2 consensus (Figure 1A). To verify whether this A/T-rich element could be recognized by cardiac-derived or recombinant MEF2 proteins, it was used in gel shift assay to compete MEF2 binding on the well characterized muscle creatine kinase (MCK) MEF2 site. As seen in Figure 1B, several A/T-rich ANF elements including the TATA-box, the SRE-like and the CArG-box could not compete the MEF2 binding on the MCK probe even when used at a 500-fold molar excess. On the other hand, the distal A/T-rich element was able to displace the MEF2 binding although at a much lower efficiency than the MCK site. Identical results were obtained using MEF2C and




Fig. 2. The low-affinity A/T-rich and the proximal GATA elements contribute to MEF2-dependent ANF promoter activation. (A) Dose-dependent ANF₇₀₀ promoter activation by MEF2A, MEF2C and MEF2D in HeLa, CV-1 and P19 cell lines. Transient transfections were performed using 50 ng, 100 ng, 500 ng and 1 μ g of MEF2 expression vector. Note the fold-activation difference between HeLa and CV-1 or P19 cells. (B) Preferential activation of the ANF₇₀₀ and α MHC promoters, but not an artificial MEF2 reporter, in HeLa cells. Transfections were performed using 1 μ g of MEF2A expression vector. Similar results were obtained using MEF2C and MEF2D. (C) The low-affinity A/T-rich and the proximal GATA elements contribute to MEF2-dependent ANF promoter activation. Transfections were performed in HeLa cells using 1 μ g of MEF2A expression vector. Similar results were obtained using MEF2C and MEF2D. 3XMEF2 and 2XA/T are the MEF2-MCK and the ANF A/T-rich elements trimerized and dimerized, respectively, in front of the ANF₅₀ minimal promoter.

MEF2D or cardiomyocyte nuclear extracts (data not shown). Consistent with its ability to recognize MEF2 proteins, the ANF A/T-rich probe was able to bind all three recombinant MEF2 proteins tested (MEF2A, MEF2C and MEF2D) albeit with low affinity (Figure 1B and data not shown). Finally, when cloned upstream of a minimal promoter, the ANF A/T-rich element could be transactivated 4-fold by cotransfection with MEF2 expression vectors in heterologous cells (Figure 2C). These results suggest that the ANF promoter contains a very low-affinity MEF2-binding site that could mediate MEF2 action.

Indeed, MEF2C and other MEF2 proteins activate the ANF promoter in a dose-dependent manner in several non-cardiac cells (Figure 2A). Interestingly, the magnitude of activation was much greater (4- to 6-fold) in HeLa cells than in most cell lines tested (including CV1, P19 and C2C12), with maximal ANF promoter induction of 15-fold. This difference in MEF2 responsiveness was also observed with the cardiac α MHC promoter, which was induced by 10-fold in HeLa cells and was barely responsive in CV1 or P19 cells (Figure 2B and data not shown); in contrast, a synthetic promoter harboring a multimerized MCK MEF2 site upstream of a minimal ANF promoter was more similarly activated by MEF2 in HeLa (9-fold) and CV1 (6-fold) cells (Figure 2B). The transfected MEF2 vectors produced similarly high levels of MEF2 proteins in all cell lines as assessed by gel shift assays (data not shown). Thus, the differences observed in the level of MEF2-dependent ANF and α MHC promoter activation may reflect cooperative interaction between




Fig. 3. The MEF2 and GATA transcription factors cooperatively activate the ANF₇₀₀ promoter. (A) MEF2A, MEF2C and MEF2D functionally interact with GATA-4. Cotransfections were performed in HeLa cells using the ANF-luc₇₀₀ construct and 1 μ g of MEF2A, MEF2C or MEF2D expression vector in the absence (-) or presence (+) of 1.5 μ g of GATA-4 expression vector. (B) MEF2 proteins functionally interact with a subset of GATA proteins. Cotransfections were performed as in (A) using 1.5 μ g of various GATA expression vectors in the absence (-) or presence (+) of 1 μ g of MEF2C expression vector. Note that cooperative interaction between MEF2A and the different GATA factors was identical to the one shown here for MEF2C and GATA-1 to -6. Similar results were also obtained in the CV1 cell line.




Fig. 4. The proximal GATA element is necessary and sufficient for MEF2-GATA synergy. Cotransfections were performed in HeLa cells using various promoter constructs and 1 μ g of MEF2A and/or 1.5 μ g of GATA-4 expression vectors. The ANF promoter constructs used are described in Materials and methods. 3XMEF2 and 2XA/T are the MEF2-MCK and the ANF A/T-rich elements trimerized and dimerized, respectively, in front of the ANF₋₅₀ minimal promoter. 2XGATA is a dimer of the BNP GATA elements in front of the minimal BNP promoter.

transfected MEF2 proteins and other cellular factors bound to the promoters.

Mutational analysis was used to test which DNA elements on the ANF promoter are required for activation by MEF2 (Figure 2C). Consistent with its characterization as a weak-affinity MEF2 site, the A/T-rich element was necessary for maximal MEF2 activation but its deletion or mutation reduced promoter activation by only 30%. Surprisingly, mutation of the proximal GATA element that can bind endogenous GATA-2 protein present in HeLa cells (Grépin *et al.*, 1994) had a more drastic effect on MEF2 responsiveness, suggesting that GATA factors may cooperate with MEF2. In fact, the proximal ANF promoter (ANF₋₁₃₅), which lacks any MEF2-binding site, was induced 4- to 5-fold by MEF2 proteins, and mutation of the GATA site therein abrogated MEF2 responsiveness (Figure 2C). This element is a high-affinity binding site for GATA factors (Charron *et al.*, 1999) but does not bind MEF2 proteins (Figure 1B). Together, these results suggest that MEF2 could act as a cofactor of promoter-bound GATA proteins to activate the ANF and possibly other cardiac promoters.

Synergistic activation of the ANF promoter by MEF2 and GATA factors

Two members of the GATA family of zinc finger transcription factors, GATA-4 and GATA-6, are expressed in cardiac muscle cells and bind to and activate the ANF promoter (Charron *et al.*, 1999). To test whether either factor could recruit MEF2 proteins to target promoters, the effect of co-expressing them with MEF2 in heterologous cells was assayed on ANF promoter activity. Cotransfection of GATA-4 with MEF2A, MEF2C or MEF2D leads to a synergistic 40- to 50-fold activation of the ANF promoter (Figure 3A). MEF2 proteins were also able to cooperate to varying degrees with other GATA factors including the hemopoietic GATA-1, -2 and -3, and the other cardiac GATA factor, GATA-6, but not GATA-5 (Figure 3B). The inability of MEF2C and MEF2A to synergize with GATA-5 and the more modest synergy achieved with GATA-2 and -3 are not due to different levels of GATA proteins produced as all expression vectors have been shown to produce similar protein levels (Viger *et al.*, 1998; Nemer *et al.*, 1999).

[illegible][illegible][illegible][illegible]

10
 11
 12
 13
 14
 15
 16
 17
 18
 19
 20
 21
 22
 23
 24
 25
 26
 27
 28
 29
 30
 31
 32
 33
 34
 35
 36
 37
 38
 39
 40
 41
 42
 43
 44
 45
 46
 47
 48
 49
 50
 51
 52
 53
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65
 66
 67
 68
 69
 70
 71
 72
 73
 74
 75
 76
 77
 78
 79
 80
 81
 82
 83
 84
 85
 86
 87
 88
 89
 90
 91
 92
 93
 94
 95
 96
 97
 98
 99
 100
 101
 102
 103
 104
 105
 106
 107
 108
 109
 110
 111
 112
 113
 114
 115
 116
 117
 118
 119
 120
 121
 122
 123
 124
 125
 126
 127
 128
 129
 130
 131
 132
 133
 134
 135
 136
 137
 138
 139
 140
 141
 142
 143
 144
 145
 146
 147
 148
 149
 150
 151
 152
 153
 154
 155
 156
 157
 158
 159
 160
 161
 162
 163
 164
 165
 166
 167
 168
 169
 170
 171
 172
 173
 174
 175
 176
 177
 178
 179
 180
 181
 182
 183
 184
 185
 186
 187
 188
 189
 190
 191
 192
 193
 194
 195
 196
 197
 198
 199
 200
 201
 202
 203
 204
 205
 206
 207
 208
 209
 210
 211
 212
 213
 214
 215
 216
 217
 218
 219
 220
 221
 222
 223
 224
 225
 226
 227
 228
 229
 230
 231
 232
 233
 234
 235
 236
 237
 238
 239
 240
 241
 242
 243
 244
 245
 246
 247
 248
 249
 250
 251
 252
 253
 254
 255
 256
 257
 258
 259
 260
 261
 262
 263
 264
 265
 266
 267
 268
 269
 270
 271
 272
 273
 274
 275
 276
 277
 278
 279
 280
 281
 282
 283
 284
 285
 286
 287
 288
 289
 290
 291
 292
 293
 294
 295
 296
 297
 298
 299
 300
 301
 302
 303
 304
 305
 306
 307
 308
 309
 310
 311
 312
 313
 314
 315
 316
 317
 318
 319
 320
 321
 322
 323
 324
 325
 326
 327
 328
 329
 330
 331
 332
 333
 334
 335
 336
 337
 338
 339
 340
 341
 342
 343
 344
 345
 346
 347
 348
 349
 350
 351
 352
 353
 354
 355
 356
 357
 358
 359
 360
 361
 362
 363
 364
 365
 366
 367
 368
 369
 370
 371
 372
 373
 374
 375
 376
 377
 378
 379
 380
 381
 382
 383
 384
 385
 386
 387
 388
 389
 390
 391
 392
 393
 394
 395
 396
 397
 398
 399
 400
 401
 402
 403
 404
 405
 406
 407
 408
 409
 410
 411
 412
 413
 414
 415
 416
 417
 418
 419
 420
 421
 422
 423
 424
 425
 426
 427
 428
 429
 430
 431
 432
 433
 434
 435
 436
 437
 438
 439
 440
 441
 442
 443
 444
 445
 446
 447
 448
 449
 450
 451
 452
 453
 454
 455
 456
 457
 458
 459
 460
 461
 462
 463
 464
 465
 466
 467
 468
 469
 470
 471
 472
 473
 474
 475
 476
 477
 478
 479
 480
 481
 482
 483
 484
 485
 486
 487
 488
 489
 490
 491
 492
 493
 494
 495
 496
 497
 498
 499
 500
 501
 502
 503
 504
 505
 506
 507
 508
 509
 510
 511
 512
 513
 514
 515
 516
 517
 518
 519
 520
 521
 522
 523
 524
 525
 526
 527
 528
 529
 530
 531
 532

[illegible][illegible]

In order to map the interaction domain between MEF2 and GATA proteins, different mutants of GATA-4 were *in vitro* cotranslated with MEF2A and co-immunoprecipitated using an antibody directed against the extreme C-terminus of GATA-4 (present in all mutants tested). MEF2A was able to interact with the full-length GATA-4 and the N-terminal activation domain-deleted mutant (201–443) (Figure 5C). Deletion of the N-terminal zinc finger of GATA-4 (242–443) reduced but did not abrogate interaction with MEF2A. However, MEF2A was unable to interact with the G4m [which harbors a point mutation in the C-terminal zinc finger, abolishing DNA binding (Charron *et al.*, 1999)] or the Δ 303–390 mutant, indicating that the C-terminal zinc finger structure and the basic region are essential for physical interaction with MEF2. The same approach was also used to map the GATA-4 interaction domain on MEF2 and revealed that the DNA-binding domain consisting of the MADS and MEF2 domains (MEF2A DIVE, aa 1–86) is sufficient for interaction with GATA-4 (Figure 5D). Interestingly, within the MADS domain, interaction with GATA factors and binding to DNA could be segregated as two DNA-binding-defective mutants (MEF2C R3T and MEF2C R24L) retained the ability to bind GATA-4 (Figure 5E). These results suggest that GATA-4 and MEF2 physically interact through their DNA-binding domains.

MEF2-GATA synergy does not require MEF2 DNA binding

To determine whether the activation domains of either or both GATA and MEF2 proteins are required, various GATA-4 mutants were tested for their capacity to activate the ANF promoter synergistically with MEF2A. The GATA-4 mutants that delete the N-terminal region (127–443 and 201–443) and the first zinc finger (242–443) were able to synergize with MEF2A (Figure 6A). However, the GATA-4 mutants that delete the C-terminal transactivation domain (1–332, 201–332 and 242–332) were all unable to support MEF2 synergy. Consistent with a requirement for GATA-4 DNA binding (Figure 4A) and the GATA-4 DNA-binding domain for physical interaction with MEF2 (Figure 5C), a point mutation in the second zinc finger that destroys DNA binding (G4m) abrogated MEF2 synergy (Figure 6A).

Functional synergy also required the activation domain of MEF2, as deletion of the C-terminal activation domain (MEF2A DIVE) completely abolished the synergy with GATA-4 (Figure 6B). However, consistent with the requirement for GATA- but not MEF2-binding sites, MEF2C mutants that are DNA-binding defective retained the ability to synergize with GATA-4 (Figure 6B). These results indicate that GATA-4 is able to recruit DNA-binding-defective MEF2 proteins to transcriptionally active complexes.

MEF2-GATA synergy: a mechanism for MEF2 action in the heart

We next tested whether transcription of other cardiac genes is cooperatively activated by MEF2 and GATA-4. As seen in Figure 7A, in addition to ANF, the α MHC, α -CA and B-type natriuretic peptide (BNP) promoters are also synergistically activated by MEF2 and GATA-4. Both




Fig. 6. Mapping of the GATA-4 and MEF2 domains required for synergy. (A) The C-terminal activation domain of GATA-4 is required for MEF2 synergy. Cotransfections were performed in HeLa cells on the ANF₇₀₀ promoter construct using 1 μ g of MEF2 and 1.5 μ g of GATA-4 expression vectors. (B) The C-terminal activation domain of MEF2, but not its DNA-binding capacity, is required for synergy with GATA-4, as shown by the ability of MEF2C R3T and MEF2C R24L to synergize with GATA-4. Note that the DNA-binding domain (MEF2A DIVE) is not sufficient to support functional synergy although it interacts physically with GATA-4 as shown in the previous figure.

α MHC and α -CA are downregulated in MEF2C null mice and neither contain a high-affinity MEF2-binding site although both are GATA targets (Sepulveda *et al.*, 1998; Charron *et al.*, 1999). However, as shown by the β MHC promoter, not all GATA target promoters are synergistically activated by MEF2, suggesting that functional GATA-MEF2 synergy is promoter context dependent and may be targeted to a specific subset of cardiac genes.

Finally, to ascertain whether, in cardiac cells, MEF2 proteins are GATA cofactors, the effect of a dominant-negative MEF2 protein on the activity of the proximal ANF promoter was determined. This promoter contains a GATA- but no MEF2-binding site. Cotransfection of a MEF2 mutant that retains the ability to associate physically with GATA-4 but lacks the activation domain reduces by 50% the activity of the ANF promoter in primary cardiomyocyte cultures; in contrast, cotransfection with wild-type MEF2A induces promoter activity by 80% (Figure 7B). Both activation by wild-type MEF2 and inhibition by its dominant-negative form are blunted by point mutation of the GATA-binding site (Figure 7B). These data are consistent with a role for MEF2 proteins as co-activators of GATA factors in cardiac muscle cells and point to a novel GATA-dependent pathway for transcriptional activation by MEF2.

Discussion

The MEF2 transcription factors are key regulators of cardiac myogenesis and morphogenesis, but the molecular




Fig. 7. The MEF2-GATA-4 synergy: a mechanism for MEF2 action in the heart. (A) The MEF2-GATA-4 synergy is not limited to the ANF promoter. HeLa cells were cotransfected with 1 μ g of MEF2A and 1.5 μ g of GATA-4 expression vectors together with various cardiac promoters. Except for the cardiac α -actin promoter that was from chicken, all other promoters used are from rat and are described in Materials and methods. TK81 is the thymidine kinase -81 bp promoter. Elements shaded in black and gray are high- and low-affinity sites, respectively, as determined by DNA-binding assays. (B) A dominant-negative MEF2 protein decreases ANF promoter activity in cardiomyocytes. Primary culture of cardiomyocytes was transfected with the wild-type ANF₁₃₅ (left panel) or GATA-mutated ANF₁₃₅ promoter (GATAp mut/ANF₁₃₅, right panel) and no (-), 50 ng (+) or 1000 ng (++) of MEF2A or a dominant-negative form of MEF2A (MEF2A DIVE). The results shown represent the mean \pm SD of two independent experiments each carried out in duplicate.

basis for their actions is poorly understood. The data presented here provide evidence that, in cardiac myocytes, MEF2 proteins are recruited by the cardiac-specific GATA transcription factors to target promoters and functionally synergize with this family of tissue-restricted zinc finger proteins. This observation is reminiscent of the cooperative interaction between MEF2 proteins and the myogenic bHLH factors in skeletal muscles, and suggests that MEF2 proteins are able to interact with and potentiate the action of other classes of cell-specific transcription factors. Given the co-expression of MEF2 and GATA factors in several cell types including smooth muscle, neuronal and T cells, the GATA-dependent MEF2 pathway described in this work may provide a molecular paradigm for understanding the mechanisms of action of MEF2 in many target cells.

GATA proteins are evolutionarily conserved cell-restricted transcription factors that play crucial roles in differentiation. In vertebrates, six GATA factors have been identified and they are all expressed in a lineage-restricted and developmentally controlled manner. GATA-1, -2 and -3 are predominantly expressed in hemopoietic cells while GATA-4, -5 and -6 are largely restricted to the heart and gut. Genetic and biochemical studies have revealed crucial roles for specific family members in hemopoietic, cardiac, neuronal and endodermal cells (Pevny *et al.*, 1991; Tsai *et al.*, 1994; Grépin *et al.*, 1995; Pandolfi *et al.*, 1995; Molkenstein *et al.*, 1997; Morrissey *et al.*, 1998). In addition to their essential roles in development, GATA factors are also required for the proper function of adult organs.

GATA-binding sites are present on many hemopoietic and cardiac promoters, which are potently activated by GATA factors (reviewed in Charron and Nemer, 1999). Moreover, GATA proteins act cooperatively with other classes of transcription factors including several zinc finger proteins, such as SP1 and FOG-1 (friend of GATA-1) and homeodomain-containing factors like the cardiac-specific Nkx2-5 (Durocher *et al.*, 1997; Tsang *et al.*, 1997). The differential interactions of GATA proteins with other transcription factors are likely to be important for functional specificity of GATA proteins during embryonic development and in differentiated and adult cells.

In this study, we report for the first time that, in addition to interacting with homeodomain- and zinc finger-containing proteins, GATA factors are also able to interact with members of the MADS-box family of transcription factors. This interaction involves physical contact between the C-terminal zinc finger DNA-binding domain and the adjacent basic region of GATA-4 and the MADS domain of MEF2. This, in turn, leads to synergistic activation of the ANF and other MEF2 target promoters independently of the DNA-binding activity of MEF2. Synergy is observed with two of the three cardiac GATA factors: GATA-4 and -6 but not with GATA-5; MEF2 synergy was also detected with GATA-1, -2 and -3 although at varying levels. Thus, all GATA factors are not equally competent to synergize with MEF2. The observation that GATA-4 and -6 but not -5 synergize with MEF2 is interesting given

that GATA-4 and -5 but not -6 synergize with the cardiac homeodomain protein Nkx2-5 (Durocher *et al.*, 1997); this suggests that while MEF2 and Nkx2-5 interact with the same domain of GATA-4, they apparently do not recognize the same molecular determinants.

Although GATA factors physically contact MEF2 proteins through the DNA-binding domain, DNA binding and GATA interaction are dissociable and neither physical nor functional interaction with GATA-4 on natural promoters requires MEF2 DNA binding. The ability of MEF2 to cooperate with MyoD in skeletal myogenesis and to synergize with MyoD in activating an artificial promoter driven by multimerized MyoD-binding sites was also shown to be independent of MEF2 DNA-binding capacity (Molkentin *et al.*, 1995). DNA binding has been shown to be dispensable for some of the actions of two other sequence-specific DNA-binding proteins, the glucocorticoid receptor (Reichardt *et al.*, 1998) and the cell-specific homeodomain protein Pit1 (Dasen *et al.*, 1999). However, in both cases, DNA-binding-independent activities involved transcriptional repression. In the present study, we show that transcriptional activation of several natural promoters by MEF2 proteins is independent of MEF2-binding sites and MEF2 DNA-binding ability. Thus, both suppressive and activating functions of transcription factors may involve DNA-binding-independent pathways.

In addition to the MADS domain, the activation domain of MEF2 is required for functional synergy with GATA-4, suggesting that MEF2 proteins potentiate transcriptional activity of GATA factors through recruitment and/or stabilization of co-activators in the GATA transcription complex. Such co-activators may include the CBP/p300 family, as both MEF2 and GATA proteins have been shown to interact independently with these co-activators (Sartorelli *et al.*, 1997; Blobel *et al.*, 1998; Kakita *et al.*, 1999). Alternatively, MEF2-GATA interaction may displace or overcome a corepressor of either or both factors. For example, MEF2 interaction with GATA-4 or -6 in the heart may displace FOG-2, a GATA-associated cofactor that represses GATA-4 activity (Lu *et al.*, 1999; Svensson *et al.*, 1999). Conversely, recruitment of MEF2 by GATA factors may displace the MEF2-associated corepressors MITR or the HDAC4 deacetylase (Miska *et al.*, 1999; Sparrow *et al.*, 1999).

In addition to cardiomyocytes, MEF2 proteins are co-expressed with members of the GATA family in several other cell types. Most notable is the presence of MEF2 proteins with GATA-6 in smooth muscle cells (Narita *et al.*, 1996; Suzuki *et al.*, 1996), and with GATA-3 in T lymphocytes (Zheng and Flavell, 1997), somites (George *et al.*, 1994) and brain (Pandolfi *et al.*, 1995). Given the demonstrated role of GATA factors in cell differentiation, the GATA-MEF2 synergy provides a general paradigm for understanding the role of MEF2 proteins as determining factors in diverse cell lineages.

Finally, it is tempting to speculate on the role of the MEF2-GATA synergy as a nuclear target of several signaling cascades including calcineurin and p38 MAP kinase. Both pathways, which are highly relevant to many human disorders such as ischemia, heart failure and inflammatory diseases, have been shown to activate MEF2 in cardiac (Kolodziejczyk *et al.*, 1999), skeletal muscle (Zetser *et al.*, 1999), neuronal (Mao *et al.*, 1999) and

T cells (Han *et al.*, 1997; Blaeser *et al.*, 2000). GATA factors have also been suggested as downstream targets of calcineurin in cardiac and skeletal muscle hypertrophy in connection with the calcineurin-activated NFAT factor (Molkentin *et al.*, 1998; Musaro *et al.*, 1999; Semsarian *et al.*, 1999). It would be interesting to test whether calcineurin or p38-dependent post-translational modifications of MEF2 or GATA proteins modulate the MEF2-GATA physical interaction and/or the resulting functional synergy.

Materials and methods

Cell cultures and transfections

HeLa cells were grown in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum (FBS). Transfections were carried out using calcium phosphate 24 h after plating. At 36 h post-transfection, cells were harvested and luciferase activity was assayed with a Berthold LB953 luminometer. The amount of reporter was kept at 1.5 µg per 35 mm dish and the total amount of DNA was kept constant (usually 7 µg). The amount of expression vector used is indicated in the figure legends. Primary cardiomyocyte cultures were prepared from 4-day-old Sprague-Dawley rats as previously described (Charron *et al.*, 1999). The results shown are the mean \pm SD of at least two independent experiments carried out in duplicate.

Plasmids

ANF-luciferase promoter constructs were cloned in the PXP-2 vector as described previously (Argentin *et al.*, 1994; Durocher *et al.*, 1996). The BNP-luc constructs were described in Grépin *et al.* (1994), the β MHC-luc and cardiac α -actin-luc reporters were described in Abdellatif *et al.* (1994). The α MHC-luc vector was kindly provided by P.M. Buttrick (Buttrick *et al.*, 1993). The construction of the various pCG-GATA-4 vectors was based on the original rat GATA-4 cDNA as previously described (Grépin *et al.*, 1994). The various deletions or point mutations of the ANF promoter and the pRSET-GATA-4 derivatives were generated as described previously (Durocher *et al.*, 1997; Charron *et al.*, 1999). ANF constructs with mutations in the GATA elements or in the NKE were previously described (Durocher *et al.*, 1996; Charron *et al.*, 1999). The A/T-rich mutation is shown in Figure 1; the SRE-like mutation replaces the TTT of the ANF-SRE by GGG thus destroying SRF binding. Heterologous promoters were generated by multimerizing the relevant oligonucleotides flanked by *Bam*HI and *Bgl*II sites upstream of the minimal (~50 bp) ANF-luciferase reporter. pCDNA-MEF2A DIVE was constructed by insertion of the *Xba*I-*Bam*HI fragment of the corresponding pCGN-MEF2A DIVE construct into the *Xba*I-*Bam*HI sites of the pCDNA-3 vector. MEF2 plasmids were kindly provided by E.N. Olson (Molkentin *et al.*, 1996a) and K. Walsh (Andres *et al.*, 1995). The MBP-GATA-4 plasmid was prepared by subcloning a *Xba*I-*Bam*HI rat GATA-4 cDNA fragment containing the entire open reading frame and 1.2 kb of 3' untranslated sequences (Grépin *et al.*, 1994) into the MBP-expressing pMalc-2 vector (New England Biolabs, Beverly, MA) cut with *Nhe*I-*Bam*HI.

Recombinant protein production

Recombinant MBP-GATA-4 was obtained according to our previously described protocol (Durocher *et al.*, 1997). Essentially, individual colonies were picked and grown in 500 ml of LB up to an OD of 0.6 at 600 nm. Induction of the recombinant protein was carried out by adding isopropyl- β -D-thiogalactopyranoside (IPTG) at a final concentration of 0.5 mM for 2 h at 37°C. The cultures were centrifuged and the bacteria were resuspended and lysed by sonication. Purification on amylose columns (New England Biolabs, Beverly, MA) was performed according to the manufacturer's instructions.

In vitro transcribed/translated 35 S-labeled MEF2 and GATA proteins were produced in rabbit reticulocyte lysates using the TNT-coupled *in vitro* transcription/translation system (Promega Corp., Madison, WI) from pCDNA-MEF2 derivatives using either T7 or Sp6 RNA polymerase.

In vitro protein-protein interactions

In vitro binding studies were performed using purified MBP-GATA-4 immobilized on an amylose-Sepharose resin (New England Biolabs) and *in vitro* transcribed/translated MEF2 proteins. Typically, 2-6 µl of

³⁵S-labeled MEF2 proteins were incubated in the presence of 300 ng of immobilized GATA-4 fusion protein in 500 µl of binding buffer [150 mM NaCl, 50 mM Tris-Cl pH 7.5, 0.3% Nonidet P-40, 1 mM dithiothreitol (DTT), 0.5 mM phenylmethylsulfonyl fluoride (PMSF) and 0.25% bovine serum albumin (BSA)] for 2 h at 4°C with agitation and then centrifuged for 2 min at 15 000 r.p.m. at room temperature. The resin was washed three times by vortexing in 500 µl of binding buffer at room temperature and three more times by vortexing in 500 µl of binding buffer without BSA. The protein complexes were released from the resin after boiling in Laemmli buffer and resolved by SDS-PAGE. Labeled proteins were visualized and quantified by autoradiography on phosphor storage plates (PhosphorImager, Molecular Dynamics).

To determine the domains of GATA-4 and MEF2 required for physical interaction, full-length GATA-4 or mutated GATA-4 plasmids were used for *in vitro* cotranscription/cotranslation with wild-type or mutant MEF2A and MEF2C. The cotranslated proteins were incubated in 500 µl of binding buffer with 1 µl of GATA-4 antibody (Santa-Cruz Biotechnology) for 2 h at 4°C with agitation and for an additional 2 h with 20 µl of protein A/G Plus-agarose added (Santa Cruz Biotechnology). Bound immunocomplexes were washed and visualized as mentioned above.

Electrophoretic mobility shift assays (EMSAs)

Three microliters of the *in vitro* translated MEF2A, MEF2C and MEF2D proteins were used for the binding reactions performed essentially as previously described for GATA binding (Charron et al., 1999) except that 100 ng of dl-dC were included in the binding reaction. Reactions were loaded on a 4% polyacrylamide gel and run at 200 V at room temperature in 0.25× Tris-borate-EDTA. The MEF2-MCK probe is as described in Molkenkin et al. (1996b). The rat ANF probes used were as follows: TATA-box, ⁻⁴⁶TCAGGGAGCTGGGGCTATAAAACGGGAGACGCC⁻¹¹; SRE-like, ⁻¹²⁴GATCCACTGATACTTTAAAGGGCATCTTCA⁻⁹⁹; CA/G, ⁻¹¹⁷GATCCTCCCGCCCTTATTGGAGCCCTGA⁻³⁹⁰; A/T-rich, ⁻⁵⁹⁷GATCCATCTACTAAAAAATATAATAGCTCTTCA⁻⁵⁶⁷.

Immunoprecipitations and immunoblots

Co-immunoprecipitations of Flag-GATA-4 and HA-MEF2A were carried out using nuclear extracts of 293T cells overexpressing the relevant proteins. Nuclear extracts were prepared as follows: five million 293T cells transfected with 15 µg of expression vectors were harvested 48 h post-transfection in ice-cold phosphate-buffered saline (PBS) containing 1 mM sodium orthovanadate and 1 mM EDTA. The cells were resuspended in hypotonic buffer (20 mM HEPES pH 7.9, 20 mM sodium fluoride, 1 mM sodium pyrophosphate, 1 mM sodium orthovanadate, 1 mM EDTA, 1 mM EGTA, 0.25 mM sodium molybdate, 10 µg/ml leupeptin, 10 µg/ml aprotinin, 10 µg/ml pepstatin, 2 mM DTT, 0.5 mM PMSF and 100 nM okadaic acid) and swollen on ice for 15 min. Twenty-five microliters of 10% NP-40 were added and the microtubes were vortexed vigorously. The nuclei were then pelleted by centrifugation at 7000 r.p.m. at 4°C. The nuclear pellet was resuspended in 50–100 µl of high salt buffer (hypotonic buffer containing 20% glycerol and 0.4% NaCl) and shaken vigorously at 4°C for 1 h. The nuclear extracts were cleared by centrifugation at 15 000 r.p.m. for 15 min at 4°C and the protein concentration was determined by the Bradford assay. Co-immunoprecipitation reactions were carried out on 50 µg of nuclear extracts using 1 µl of 12CA5 antibody in 500 µl of binding buffer without BSA, and bound immunocomplexes were washed and subjected to SDS-PAGE, as described previously (Durocher et al., 1997). Proteins were transferred on Hybond-PVDF membrane and subjected to immunoblotting. Anti-Flag M5 (Sigma) and 12CA5 (anti-HA) monoclonal antibodies were used at a dilution of 1/8000, revealed with an anti-mouse-HRP (Sigma) at a dilution of 1/50 000 and visualized using ECL Plus (Amersham Pharmacia Biotechnology).

Acknowledgements

We are grateful to Anne Aries and Pierre Paradis for generating some ANF mutants, and to Eric Olson, Jeffrey Molkenkin and Ken Walsh for the gift of MEF2 vectors. We thank Lise Larocque for secretarial assistance, and members of the Nemer laboratory for discussions and critical reading of this manuscript. This work was supported by grants from the Canadian Medical Research Council (MRC). S.M. holds a studentship from the MRC; F.C. was funded by a research traineeship from the Heart and Stroke Foundation of Canada and is presently a recipient of a National Cancer Institute of Canada studentship. M.N. is a senior MRC scientist.

References

- Abdellatif, M., MacLellan, W.R. and Schneider, M.D. (1994) p21 Ras as a governor of global gene expression. *J. Biol. Chem.*, **269**, 15423–15426.
- Andres, V., Fisher, S., Wearsch, P. and Walsh, K. (1995) Regulation of GAX homeobox gene transcription by a combination of positive factors including myocyte-specific enhancer factor 2. *Mol. Cell. Biol.*, **15**, 4272–4281.
- Argentin, S., Ardati, A., Tremblay, S., Lihmann, J., Robitaille, L., Drouin, J. and Nemer, M. (1994) Developmental stage-specific regulation of atrial natriuretic factor gene transcription in cardiac cells. *Mol. Cell. Biol.*, **14**, 777–790.
- Bi, W., Drake, C.J. and Schwarz, J.J. (1999) The transcription factor MEF2C-null mouse exhibits complex vascular malformations and reduced cardiac expression of angiopoietin 1 and VEGF. *Dev. Biol.*, **211**, 255–267.
- Black, B.L. and Olson, E.N. (1998) Transcriptional control of muscle development by myocyte enhancer factor-2 (MEF2) proteins. *Annu. Rev. Cell. Dev. Biol.*, **14**, 167–196.
- Black, B.L., Molkenkin, J.D. and Olson, E.N. (1998) Multiple roles for the MyoD basic region in transmission of transcriptional activation signals and interaction with MEF2. *Mol. Cell. Biol.*, **18**, 69–77.
- Blaeser, F., Ho, N., Prywes, R. and Chatila, T.A. (2000) Ca²⁺-dependent gene expression mediated by MEF2 transcription factors. *J. Biol. Chem.*, **275**, 197–209.
- Blobel, G.A., Nakajima, T., Eckner, R., Montminy, M. and Orkin, S.H. (1998) CREB-binding protein cooperates with transcription factor GATA-1 and is required for erythroid differentiation. *Proc. Natl. Acad. Sci. USA*, **95**, 2061–2066.
- Bour, B.A., O'Brien, M.A., Lockwood, W.L., Goldstein, E.S., Bodmer, R., Taghert, P.H., Abmayr, S.M. and Nguyen, H.T. (1995) *Drosophila* MEF2, a transcription factor that is essential for myogenesis. *Genes Dev.*, **9**, 730–741.
- Buttrick, P.M., Kaplan, M.L., Kitsis, R.N. and Leinwand, L.A. (1993) Distinct behavior of cardiac myosin heavy chain gene constructs *in vivo*. Discordance with *in vitro* results. *Circ. Res.*, **72**, 1211–1217.
- Chambers, A.E., Logan, M., Kotecha, S., Towers, N., Sparrow, D. and Mohun, T.J. (1994) The RSRF/MEF2 protein SL1 regulates cardiac muscle-specific transcription of a myosin light-chain gene in *Xenopus* embryos. *Genes Dev.*, **8**, 1324–1334.
- Charron, F. and Nemer, M. (1999) GATA transcription factors and cardiac development. *Semin. Cell Dev. Biol.*, **10**, 85–91.
- Charron, F., Paradis, P., Bronchain, O., Nemer, G. and Nemer, M. (1999) Cooperative interaction between GATA-4 and GATA-6 regulates myocardial gene expression. *Mol. Cell. Biol.*, **19**, 4355–4365.
- Dasen, J.S., O'Connell, S.M., Flynn, S.E., Treier, M., Gleiberman, A.S., Szeto, D.P., Hooshmand, F., Aggarwal, A.K. and Rosenfeld, M.G. (1999) Reciprocal interactions of Pit1 and GATA2 mediate signaling gradient-induced determination of pituitary cell types. *Cell*, **97**, 587–598.
- Di Lisi, R., Millino, C., Calabria, E., Altruda, F., Schiaffino, S. and Ausoni, S. (1998) Combinatorial *cis*-acting elements control tissue-specific activation of the cardiac troponin I gene *in vitro* and *in vivo*. *J. Biol. Chem.*, **273**, 25371–25380.
- Durocher, D., Chen, C.Y., Ardati, A., Schwartz, R.J. and Nemer, M. (1996) The ANF promoter is a downstream target for Nkx-2.5 in the myocardium. *Mol. Cell. Biol.*, **16**, 4648–4655.
- Durocher, D., Charron, F., Warren, R., Schwartz, R.J. and Nemer, M. (1997) The cardiac transcription factors Nkx2-5 and GATA-4 are mutual cofactors. *EMBO J.*, **16**, 5687–5696.
- Durocher, D., Grépin, C. and Nemer, M. (1998) Regulation of gene expression in the endocrine heart. In Conn, P.M. (ed.), *Recent Progress in Hormone Research*. The Endocrine Society Press, Bethesda, MD, pp. 7–23.
- Fu, Y.C. and Izumo, S. (1995) Cardiac myogenesis—overexpression of *xcsx2* or *xmf2a* in whole *Xenopus* embryos induces the precocious expression of *xmhc-α* gene. *Roux Arch. Dev. Biol.*, **205**, 198–202.
- George, K.M., Leonard, M.W., Roth, M.E., Lieuw, K.H., Kioussis, D., Grosfeld, F. and Engel, J.D. (1994) Embryonic expression and cloning of the murine GATA-3 gene. *Development*, **120**, 2673–2686.
- Grépin, C., Dagnino, L., Robitaille, L., Haberstroh, L., Antakly, T. and Nemer, M. (1994) A hormone-encoding gene identifies a pathway for cardiac but not skeletal muscle gene transcription. *Mol. Cell. Biol.*, **14**, 3115–3129.
- Grépin, C., Robitaille, L., Antakly, T. and Nemer, M. (1995) Inhibition of

- transcription factor GATA-4 expression blocks *in vitro* cardiac muscle differentiation. *Mol. Cell. Biol.*, 15, 4095-4102.
- Han, J., Jiang, Y., Li, Z., Kravchenko, V.V. and Ulevitch, R.J. (1997) Activation of the transcription factor MEF2C by the MAP kinase p38 in inflammation. *Nature*, 386, 296-299.
- Iannello, R.C., Mar, J.H. and Ordahl, C.P. (1991) Characterization of a promoter element required for transcription in myocardial cells. *J. Biol. Chem.*, 266, 3309-3316.
- Kakita, T., Hasegawa, K., Morimoto, T., Kaburagi, S., Wada, H. and Sasayama, S. (1999) p300 protein as a coactivator of GATA-5 in the transcription of cardiac-restricted atrial natriuretic factor gene. *J. Biol. Chem.*, 274, 34096-34102.
- Kaushal, S., Schneider, J.W., Nadal-Ginard, B. and Mahdavi, V. (1994) Activation of the myogenic lineage by *mef2a*, a factor that induces and cooperates with *myod*. *Science*, 266, 1236-1240.
- Kolodziejczyk, S.M., Wang, L., Balazsi, K., DeRepentigny, Y., Kothary, R. and Megeney, L.A. (1999) MEF2 is upregulated during cardiac hypertrophy and is required for normal post-natal growth of the myocardium. *Curr. Biol.*, 9, 1203-1206.
- Kuisk, I.R., Li, H., Tran, D. and Capetanaki, Y. (1996) A single MEF2 site governs desmin transcription in both heart and skeletal muscle during mouse embryogenesis. *Dev. Biol.*, 174, 1-13.
- Lee, Y., Nadal-Ginard, B., Mahdavi, V. and Izumo, S. (1997) Myocyte-specific enhancer factor 2 and thyroid hormone receptor associate and synergistically activate the α -cardiac myosin heavy-chain gene. *Mol. Cell. Biol.*, 17, 2745-2755.
- Lilly, B., Zhao, B., Ranganayakulu, G., Paterson, B.M., Schulz, R.A. and Olson, E.N. (1995) Requirement of MADS domain transcription factor D-MEF2 for muscle formation in *Drosophila*. *Science*, 267, 688-693.
- Lin, Q., Schwarz, J., Bucana, C. and Olson, E.N. (1997) Control of mouse cardiac morphogenesis and myogenesis by transcription factor *mef2c*. *Science*, 276, 1404-1407.
- Lin, Q., Lu, J., Yanagisawa, H., Webb, R., Lyons, G.E., Richardson, J.A. and Olson, E.N. (1998) Requirement of the MADS-box transcription factor MEF2C for vascular development. *Development*, 125, 4565-4574.
- Lu, J.R., McKinsey, T.A., Xu, H.T., Wang, D.Z., Richardson, J.A. and Olson, E.N. (1999) FOG-2, a heart- and brain-enriched cofactor for GATA transcription factors. *Mol. Cell. Biol.*, 19, 4495-4502.
- Mao, Z., Bonni, A., Xia, F., Nadal-Vicens, M. and Greenberg, M.E. (1999) Neuronal activity-dependent cell survival mediated by transcription factor MEF2. *Science*, 286, 785-790.
- Miska, E.A., Karlsson, C., Langley, E., Nielsen, S.J., Pines, J. and Kouzarides, T. (1999) HDAC4 deacetylase associates with and represses the MEF2 transcription factor. *EMBO J.*, 18, 5099-5107.
- Molkentin, J.D. and Markham, B.E. (1993) Myocyte-specific enhancer-binding factor (MEF-2) regulates α -cardiac myosin heavy chain gene expression *in vitro* and *in vivo*. *J. Biol. Chem.*, 268, 19512-19520.
- Molkentin, J.D., Black, B.L., Martin, J.F. and Olson, E.N. (1995) Cooperative activation of muscle gene expression by MEF2 and myogenic bHLH proteins. *Cell*, 83, 1125-1136.
- Molkentin, J.D., Firulli, A.B., Black, B.L., Martin, J.F., Hustad, C.M., Copeland, N., Jenkins, N., Lyons, G. and Olson, E.N. (1996a) MEF2B is a potent transactivator expressed in early myogenic lineages. *Mol. Cell. Biol.*, 16, 3814-3824.
- Molkentin, J.D., Black, B.L., Martin, J.F. and Olson, E.N. (1996b) Mutational analysis of the DNA binding, dimerization, and transcriptional activation domains of MEF2C. *Mol. Cell. Biol.*, 16, 2627-2636.
- Molkentin, J.D., Lin, Q., Duncan, S.A. and Olson, E.N. (1997) Requirement of the transcription factor GATA4 for heart tube formation and ventral morphogenesis. *Genes Dev.*, 11, 1061-1072.
- Molkentin, J.D., Lu, J.R., Antos, C.L., Markham, B., Richardson, J., Robbins, J., Grant, S.R. and Olson, E.N. (1998) A calcineurin-dependent transcriptional pathway for cardiac hypertrophy. *Cell*, 93, 215-228.
- Morrissey, E.E., Tang, Z., Sigrist, K., Lu, M.M., Jiang, F., Ip, H.S. and Parmacek, M.S. (1998) GATA6 regulates HNF4 and is required for differentiation of visceral endoderm in the mouse embryo. *Genes Dev.*, 12, 3579-3590.
- Musaro, A., McCullagh, K.J., Naya, F.J., Olson, E.N. and Rosenthal, N. (1999) IGF-1 induces skeletal myocyte hypertrophy through calcineurin in association with GATA-2 and NF-ATc1. *Nature*, 400, 581-585.
- Narita, N., Heikinheimo, M., Bielinska, M., White, R.A. and Wilson, D.B. (1996) The gene for transcription factor GATA-6 resides on mouse chromosome 18 and is expressed in myocardium and vascular smooth muscle. *Genomics*, 36, 345-348.
- Nemer, G., Qureshi, S.A., Malo, D. and Nemer, M. (1999) Functional analysis and chromosomal mapping of GATA5, a gene encoding a zinc finger DNA-binding protein. *Mamm. Genome*, 10, 993-999.
- Pandolfi, P.P., Roth, M.E., Karis, A., Leonard, M.W., Dzierzak, E., Grosfeld, F.G., Engel, J.D. and Lindenbaum, M.H. (1995) Targeted disruption of the GATA3 gene causes severe abnormalities in the nervous system and in fetal liver haematopoiesis. *Nature Genet.*, 11, 40-44.
- Pevny, L., Simon, M.C., Robertson, E., Klein, W.H., Tsai, S.F., D'Agati, V., Orkin, S.H. and Costantini, F. (1991) Erythroid differentiation in chimaeric mice blocked by a targeted mutation in the gene for transcription factor GATA-1. *Nature*, 349, 257-260.
- Ranganayakulu, G., Zhao, B., Dokidis, A., Molkentin, J.D., Olson, E.N. and Schulz, R.A. (1995) A series of mutations in the D-MEF2 transcription factor reveal multiple functions in larval and adult myogenesis in *Drosophila*. *Dev. Biol.*, 171, 169-181.
- Reichardt, H.M. et al. (1998) DNA binding of the glucocorticoid receptor is not essential for survival. *Cell*, 93, 531-541.
- Sartorelli, V., Huang, J., Hamamori, Y. and Kedes, L. (1997) Molecular mechanisms of myogenic coactivation by p300: direct interaction with the activation domain of MyoD and with the MADS box of MEF2C. *Mol. Cell. Biol.*, 17, 1010-1026.
- Semsarian, C., Wu, M.J., Ju, Y.K., Marciniak, T., Yeoh, T., Allen, D.G., Harvey, R.P. and Graham, R.M. (1999) Skeletal muscle hypertrophy is mediated by a Ca^{2+} -dependent calcineurin signalling pathway. *Nature*, 400, 576-581.
- Sepulveda, J.L., Belaguli, N., Nigam, V., Chen, C.Y., Nemer, M. and Schwartz, R.J. (1998) GATA-4 and Nkx-2.5 coactivate Nkx-2 DNA binding targets: role for regulating early cardiac gene expression. *Mol. Cell. Biol.*, 18, 3405-3415.
- Sparrow, D.B., Miska, E.A., Langley, E., Reynaud-Deonauth, S., Kotecha, S., Towers, N., Spohr, G., Kouzarides, T. and Mohun, T.J. (1999) MEF-2 function is modified by a novel co-repressor, MITR. *EMBO J.*, 18, 5085-5098.
- Suzuki, E., Evans, T., Lowry, J., Truong, L., Bell, D.W., Testa, J.R. and Walsh, K. (1996) The human GATA-6 gene: structure, chromosomal location, and regulation of expression by tissue-specific and mitogen-responsive signals. *Genomics*, 38, 283-290.
- Svensson, E.C., Tufts, R.L., Polk, C.E. and Leiden, J.M. (1999) Molecular cloning of FOG-2: a modulator of transcription factor GATA-4 in cardiomyocytes. *Proc. Natl Acad. Sci. USA*, 96, 956-961.
- Tsai, F.Y., Keller, G., Kuo, F.C., Weiss, M., Chen, J., Rosenblatt, M., Alt, F.W. and Orkin, S.H. (1994) An early haematopoietic defect in mice lacking the transcription factor GATA-2. *Nature*, 371, 221-226.
- Tsang, A.P., Visvader, J.E., Turner, C.A., Fujiwara, Y., Yu, C., Weiss, M.J., Crossley, M. and Orkin, S.H. (1997) FOG, a multitype zinc finger protein, acts as a cofactor for transcription factor GATA-1 in erythroid and megakaryocytic differentiation. *Cell*, 90, 109-119.
- Viger, R.S., Mertineit, C., Trasler, J.M. and Nemer, M. (1998) Transcription factor GATA-4 is expressed in a sexually dimorphic pattern during mouse gonadal development and is a potent activator of the Müllerian inhibiting substance promoter. *Development*, 125, 2665-2675.
- Yu, Y.T., Breitbart, R.E., Smoot, L.B., Lee, Y., Mahdavi, V. and Nadal-Ginard, B. (1992) Human myocyte-specific enhancer factor 2 comprises a group of tissue-restricted MADS box transcription factors. *Genes Dev.*, 6, 1783-1798.
- Zetser, A., Gredinger, E. and Bengal, E. (1999) p38 mitogen-activated protein kinase pathway promotes skeletal muscle differentiation. Participation of the *Mef2c* transcription factor. *J. Biol. Chem.*, 274, 5193-5200.
- Zheng, W. and Flavell, R.A. (1997) The transcription factor GATA-3 is necessary and sufficient for Th2 cytokine gene expression in CD4 T cells. *Cell*, 89, 587-596.
- Zhu, H., Garcia, A.V., Ross, R.S., Evans, S.M. and Chien, K.R. (1991) A conserved 28-base-pair element (HF-1) in the rat cardiac myosin light-chain-2 gene confers cardiac-specific and α -adrenergic-inducible expression in cultured neonatal rat myocardial cells. *Mol. Cell. Biol.*, 11, 2273-2281.

Received January 27, 2000; revised and accepted March 13, 2000